

Submitted to "IEEE/ACM Transactions on Networking" in August 2002

Correction to Low and Lapsley's article "Optimization Flow Control, I: Basic Algorithm and Convergence"

Andrzej Karbowski

NASK (Research and Academic Computer Network),

ul. Wwozowa 18,

02-796 Warsaw, Poland,

E-mail: A.Karbowski@ia.pw.edu.pl

Abstract

In the note an error in Low and Lapsley's article [1] is pointed out. Because of this error the proof of the Theorem 2 presented in the article is incomplete and some assessments are wrong. In the second part of the note the author proposes a correction to this proof.

1 Error in the proof of the Theorem 2

The last passage in the assessment (29) on the page 873 (the proof of Lemma 6), namely

$$\begin{aligned}
 D(p(t+1)) &\leq D(p(t)) - \left(\frac{1}{\gamma} - A_1\right) \|\pi(t)\|^2 \\
 &\quad + A_2 \sum_{t'=t-2t_0}^{t-1} \|\pi(t')\| \cdot \|\pi(t)\| \\
 &\leq D(p(t)) - \left(\frac{1}{\gamma} - A_1\right) \|\pi(t)\|^2 \\
 &\quad + A_2 \sum_{t'=t-2t_0}^t \|\pi(t)\|^2
 \end{aligned} \tag{1}$$

is incorrect. Low and Lapsley justify it: *"where the last inequality holds because the convex function $\sum_i y_i^2 + z^2 - \sum_i y_i z$ attains a unique minimum over $\{(y_i, z) \mid y_i \geq 0, z \geq 0\}$ at the origin."*

But this is not true. It is sufficient to take $\dim y = 5$ and $y = [5, 4, 3, 4, 5]$, $z = 10$. The considered function takes the value -19 .

It is so, because the Hessian of this function has the form (taking: $x = [y_1, y_2, \dots, z]'$):

$$H = \begin{bmatrix} 2 & 0 & 0 & \dots & 0 & -1 \\ 0 & 2 & 0 & \dots & 0 & -1 \\ 0 & 0 & 2 & \dots & 0 & -1 \\ \vdots & & & & & \\ 0 & 0 & 0 & \dots & 2 & -1 \\ -1 & -1 & -1 & \dots & -1 & 2 \end{bmatrix} \tag{2}$$

and its characteristic polynomial (e.g. calculated from the Schur's formula: for $A : n \times n$; $D : m \times m$, $B^T, C : m \times n$; $\det([A, B; C, D]) = \det(A) \cdot \det(D - CA^{-1}B)$) will be:

$$\det(H - \lambda I) = (2 - \lambda)^{n-1}(\lambda^2 - 4\lambda + 4 - n) \tag{3}$$

where $n = \dim y$. In this way we will have eigenvalues: $\lambda_i = 2, i = 1, \dots, n-1$ and $\lambda_{n,n+1} = 2 \pm \sqrt{n}$. This means, that for $n > 4$ there will be one negative eigenvalue and the function will not be convex.

2 Correction to the proof of the Theorem 2

The mentioned passage in the assessment (29) in [1] (here ineq. (1)) should be changed. The changes are based on a very simple assessment:

$$\forall y_i, z \in \mathbb{R} \quad (y_i - z)^2 = y_i^2 - 2y_i z + z^2 \geq 0 \quad (4)$$

From which after elementary operations we get:

$$y_i z \leq \frac{y_i^2 + z^2}{2} \quad (5)$$

Let us apply the assessment (5) to all elements of a finite set of real numbers $\{y_i, i \in I\}$, where I is a set of integer indices, and sum up both sides of these inequalities over all $i \in I$. We will obtain:

$$\sum_{i \in I} y_i z \leq \frac{1}{2} \sum_{i \in I} (y_i^2 + z^2) = \frac{1}{2} \bar{I} \cdot z^2 + \frac{1}{2} \sum_{i \in I} y_i^2 \quad (6)$$

That is:

$$\sum_{i \in I} y_i z \leq \frac{1}{2} \bar{I} \cdot z^2 + \frac{1}{2} \sum_{i \in I} y_i^2 \quad (7)$$

where $\bar{I}$ is the number of elements of the set I . We will use the assessment (7) to transform the first part of assessment (1). In particular, owing to (7), for the last component of the right hand side we will have:

$$\sum_{t'=t-2t_0}^{t-1} \|\pi(t')\| \cdot \|\pi(t)\| \leq \frac{1}{2} [t-1 - (t-2t_0) + 1] \cdot \|\pi(t)\|^2 + \frac{1}{2} \sum_{t'=t-2t_0}^{t-1} \|\pi(t')\|^2 =$$

$$= t_0 \|\pi(t)\|^2 + \frac{1}{2} \sum_{t'=t-2t_0}^{t-1} \|\pi(t')\|^2 \quad (8)$$

Let us notice that:

$$\begin{aligned} t_0 \|\pi(t)\|^2 + \frac{1}{2} \sum_{t'=t-2t_0}^{t-1} \|\pi(t')\|^2 &= t_0 \|\pi(t)\|^2 - \frac{1}{2} \|\pi(t)\|^2 + \frac{1}{2} \|\pi(t)\|^2 + \\ &+ \frac{1}{2} \sum_{t'=t-2t_0}^{t-1} \|\pi(t')\|^2 = \left(t_0 - \frac{1}{2}\right) \|\pi(t)\|^2 + \frac{1}{2} \sum_{t'=t-2t_0}^t \|\pi(t')\|^2 \end{aligned} \quad (9)$$

So, for the value of $\sum_{t'=t-2t_0}^{t-1} \|\pi(t')\| \cdot \|\pi(t)\|$ we will have the following assessment:

$$\sum_{t'=t-2t_0}^{t-1} \|\pi(t')\| \cdot \|\pi(t)\| \leq \left(t_0 - \frac{1}{2}\right) \|\pi(t)\|^2 + \frac{1}{2} \sum_{t'=t-2t_0}^t \|\pi(t')\|^2 \quad (10)$$

The correct form of the assessment (29) in the article [1] will be then:

$$\begin{aligned} D(p(t+1)) &\leq D(p(t)) - \left(\frac{1}{\gamma} - A_1\right) \|\pi(t)\|^2 \\ &+ A_2 \sum_{t'=t-2t_0}^{t-1} \|\pi(t')\| \|\pi(t)\| \\ &\leq D(p(t)) - \left(\frac{1}{\gamma} - A_1\right) \|\pi(t)\|^2 \\ &+ A_2 \left(t_0 - \frac{1}{2}\right) \|\pi(t)\|^2 + \frac{A_2}{2} \sum_{t'=t-2t_0}^t \|\pi(t')\|^2 = \\ &= D(p(t)) - \left[\frac{1}{\gamma} - A_1 - A_2 \left(t_0 - \frac{1}{2}\right)\right] \|\pi(t)\|^2 + \frac{A_2}{2} \sum_{t'=t-2t_0}^t \|\pi(t')\|^2 \end{aligned}$$

That is:

$$D(p(t+1)) \leq D(p(t)) - \left[\frac{1}{\gamma} - A_1 - A_2 \left(t_0 - \frac{1}{2}\right)\right] \|\pi(t)\|^2 + \frac{A_2}{2} \sum_{t'=t-2t_0}^t \|\pi(t')\|^2 \quad (11)$$

Now, applying this inequality recursively to all $D(p(\tau))$, $\tau = t, t-1, \dots, 1$, taking $\pi(t) = 0$ for $t < 0$, we will get the following total assessment which replaces wrong assessment (30) in the article [1]:

$$D(p(t+1)) \leq D(p(0)) - \left[\frac{1}{\gamma} - A_1 - A_2 \left(t_0 - \frac{1}{2} \right) \right] \sum_{\tau=0}^t \|\pi(\tau)\|^2 + \frac{A_2}{2} \sum_{\tau=0}^t \sum_{t'=\tau-2t_0}^{\tau} \|\pi(t')\|^2 \quad (12)$$

Let us notice, that under the assumption that $\pi(t) = 0$ for $t < 0$:

$$\begin{aligned} \sum_{\tau=0}^t \sum_{t'=\tau-2t_0}^{\tau} \|\pi(t')\|^2 &= \sum_{\tau=0}^t \sum_{t''=0}^{2t_0} \|\pi(t'' + \tau - 2t_0)\|^2 = \sum_{t''=0}^{2t_0} \sum_{\tau=0}^t \|\pi(\tau + t'' - 2t_0)\|^2 \\ &\leq \sum_{t''=0}^{2t_0} \sum_{\tau=0}^t \|\pi(\tau)\|^2 = (2t_0 + 1) \sum_{\tau=0}^t \|\pi(\tau)\|^2 \end{aligned} \quad (13)$$

Putting (13) into (12) we finally get:

$$\begin{aligned} D(p(t+1)) &\leq D(p(0)) - \left[\frac{1}{\gamma} - A_1 - A_2 \left(t_0 - \frac{1}{2} \right) \right] \sum_{\tau=0}^t \|\pi(\tau)\|^2 + \frac{A_2}{2} (2t_0 + 1) \sum_{\tau=0}^t \|\pi(\tau)\|^2 \\ &= D(p(0)) - \left[\frac{1}{\gamma} - A_1 - A_2 \left(t_0 - \frac{1}{2} \right) - A_2 \left(t_0 + \frac{1}{2} \right) \right] \sum_{\tau=0}^t \|\pi(\tau)\|^2 = \\ &D(p(0)) - \left(\frac{1}{\gamma} - A_1 - 2A_2 t_0 \right) \sum_{\tau=0}^t \|\pi(\tau)\|^2 \end{aligned} \quad (14)$$

That is:

$$D(p(t+1)) \leq D(p(0)) - \left(\frac{1}{\gamma} - A_1 - 2A_2 t_0 \right) \sum_{\tau=0}^t \|\pi(\tau)\|^2 \quad (15)$$

References

- [1] Low, S. and D.E. Lapsley, Optimization Flow Control, I: Basic Algorithm and Convergence, *IEEE/ACM Transactions on Networking*, 7(6), pp. 861-874, 1999.